

## Chemical composition of biotite as a guide to petrogenesis of granitic rocks from Maherabad, Dehnow, Gheshlagh, Khajehmourad and Najmabad, Iran

M. H. Karimpour<sup>\*1</sup>, C. R. Stern<sup>2</sup>, M. Mouradi<sup>1</sup>

1. Research Center for Ore Deposits of Eastern Iran, Ferdowsi University of Mashhad

2. Dept. of Geological Sciences, University of Colorado, CB-399, Boulder, CO, USA

(Received: 29/11/2009, in revised form: 30/5/2010)

**Abstract:** Biotite, the dominant ferromagnesian mineral in granitoid rocks, can be used to discriminate tectonic setting, magma types and magnetite-ilmenite series. In this study, we analyzed biotite with an electron microprobe (wavelength dispersion) from different granitoids. Intrusive rocks from Maherabad porphyry Cu-Au prospecting are meta-aluminous. Biotite from Maherabad are Mg-rich type and the ratio of Fe/(Fe+Mg) is 0.286-0.309. Maherabad biotite compositions fall in the field of (sub-alkaline) calc-alkaline orogenic suites. Based on High TiO<sub>2</sub> and low Al<sub>2</sub>O<sub>3</sub> in biotites, Maherabad also belongs to magnetite series. Intrusive rocks from Najmabad, Dehnow, Gheshlagh and Khajehmourad (NDGK) are classified as belonging to the ilmenite-series of reduced S-type granitoids. Biotite from NDGK areas are Fe-rich and the Fe/(Fe+Mg) ratio in Najmabad is 0.491-0.511, in Dehnow-Kuhsangi 0.583-0.675, Gheshlagh 0.56-0.58, and Khajehmourad 0.705-0.720. respectively NDGK biotite compositions fall in the field of peraluminous granite (P) suites (S-type). Based on low TiO<sub>2</sub> and high Al<sub>2</sub>O<sub>3</sub> in biotites, Najmabad, Dehnow and Gheshlagh biotite belong to ilmenite series.

**Keywords:** Biotite, Najmabad, Maherabad, Khajehmourad, susceptibility.