

Vol.21, No. 3, Fall 1392/2013

Pages 59 to 68



# A coordination polymer of Mn (III) with pyridine-2, 3-dicarboxylic acid and 4,4'bipyridine

## J. Soleimannejad<sup>\*</sup>, E. Nazarnia

Department of Chemistry, Faculty of Sciences, Ilam University, Ilam, Iran

(Received: 4/8/2012, in revised form: 11/10/2012)

**Abstract:** The polymeric compound,  $\{(C_{10}H_9N_2)[Mn(C_7H_3NO_4)_2].(C_{10}H_8N_2).6H_2O\}_n$ , was prepared by the reaction of manganese (II) nitrate with pyridine-2,3-dicarboxylic acid, (py-2,3-dcH<sub>2</sub>) and 4,4'-bipyridine (4,4'-bipy) in a 1:2:4 molar ratio. The crystal system of the complex is monoclinic with space group P2<sub>1</sub>/n and four molecules per unit cell. The unit cell parameters are: a = 22.2099(15) Å, b = 6.6599(4) Å, c = 23.5921(16) Å and  $\beta$  = 104.014(3)°. The final R value was 0.04 based on 8960 reflections. The central atom is attached to two (py-2,3-dc)<sup>2-</sup> ligands through their N and O atoms in a planar square arrangement and two O atoms from bridging carboxylate groups, forming polymeric chains. So, the anionic complex features six-coordinate Mn (III) with a distorted octahedral geometry. In the crystal structure, a wide range of non-covalent interactions consisting of hydrogen bonding (of the types O–H···O, O–H···N and C–H···O), ion pairing, and  $\pi$ - $\pi$  (centroid-centroid distances of 3.607 and 3.721 Å), C–H··· $\pi$  (with H··· $\pi$  distance of 2.941 Å) and C=O··· $\pi$  (with O··· $\pi$  distances of 2.943, 3.702 and 3.743 Å) stacking interactions stabilize the supramolecular structure.

Keywords: Mn (III) complex; 4, 4'-bipyridine; crystal structure; hydrogen bonding.

## Introduction

Metal-based coordination polymers have been extensively studied due to their intriguing topological structures [1–3] and unique functions, such as ion-exchange, adsorption, separation, sensor and molecular recognition [4–7]. In this regard, the rational design and construction of coordination polymers with unique structural motifs and unique chemical and physical properties have attracted extensive interest in supramolecular chemistry and materials chemistry [8–10]. The framework structure of the coordination polymers is primarily dependent upon the coordination preferences of the central metal ions and the functionality of the ligands. Aside from the coordination bonding interactions, the hydrogen bonding and  $\pi$ - $\pi$  stacking interactions, the solvent molecules, counter-ions and the ratio of metal salt to organic ligand also influence the formation of the ultimate architectures.

The pyridinedicarboxylate ligands tend to be influenced by  $\pi$ - $\pi$  interaction due to the presence of aromatic pyridine ring, favoring the formation of the orderly arrangement of the molecules. It is well known that carboxylate ligands play an important role in coordination chemistry and can adopt various binding modes such as terminal monodentate, chelating to one metal center, bridging bidentate to two metal centers, and bridging tridentate to two metal centers [11].

\* Corresponding author, Tel: (021) 66495291, Fax: (021) 66495291, Email: janet\_soleimannejad@khayam.ut.ac.ir janet\_soleimannejad@yahoo.com

Among pyridinedicarboxylic acids,  $py-2,3-dcH_2$  is a good ligand for coordination and organic– inorganic hybrid compounds because of its flexible and varied coordination modes.

The rigid rod-like spacer molecules like 4,4'bipyridine (4,4'-bipy) have been widely used as building blocks for the construction of metalorganic coordination frameworks [12]. The use of organic spacers, particularly flexible dicarboxylates bridging ligands and rigid planar bidentate diimines (4,4'-bipy) as building blocks to construct various metal assemblies is of growing interest in the field of molecular materials [11]. Recently, great success has been achieved by the combination of flexible aliphatic dicarboxylates and bipy as chelating bridging ligands, which resulted in compounds having 1-3D frameworks and unique physicochemical properties [13–15]. In this pyridine-2,3-dicarboxylic study. acids molecules (py-2,3-dcH<sub>2</sub>) is coordinated to Mn(III) ions in presence of 4,4'-bipyridine (4,4'-bpy) and produced an anionic complex, this complex selfassembles with 4,4'-bipyridine (4,4'-bpy), protonated 4,4'-bipyridine (4,4'-bpyH<sup>+</sup>) and water molecules affording a compound with a polymeric structural motif.

#### **Preparation and experiments:**

A solution of Mn (NO<sub>3</sub>)<sub>2</sub>.6H<sub>2</sub>O (143 mg, 0.5

mmol), pyridine-2,3-dicarboxylic acid (167 mg, 1 mmol) and 4,4'-bipyridine (312 mg, 2 mmol) in water (10 ml) was placed in autoclave and heated up to 120°C for 2h (heating ramp 60°C/h and cooling ramp 15°C/h). Yellow crystals of the title polymeric compound were obtained after two weeks (yield 82%). Scheme 1 presents the chemical structure of the compound.

The x-ray data was collected on a Bruker SMART 1000 diffractometer (MoK $_{\alpha}$  radiation,  $\lambda = 0.71073$  Å) at 296 K.

#### **Results and discussion**

The molecular structure of the title polymeric compound is given in Figure 1. The crystal data and experimental parameters are depicted in Table 1. The crystal was solved by direct methods (SHELXS-97) and a refinement was carried out with full-matrix least-squares methods based on  $F^2$  with SHELXL-97 [16] and the final atomic coordinates of non-hydrogen atoms and selected bond lengths and bond angles are presented in Tables 2 and 3, respectively. Also hydrogen bond geometries are given separately in Table 4. This compound crystallized in the monoclinic system, space group  $P2_1/n$  with four formula units in the unit cell. The final R value was 0.04 for 8960 measured reflections.



Scheme 1



**Fig. 1** The molecular structure of the title compound, displacement ellipsoids are drawn at 50% probability level. Uncoordinated water molecules are omitted for clarity.

Table 1. Crystal and experimental	data of the title compound
$Formula = C_{34}H_{35}MnN_6O_{14}$	
Formula weight = $806.62$	
Crystal system = Monoclinic	T = 296 (2) K
Space group = $P2_{l}/n$	Z = 4
a = 22.2099(15)  Å	
b = 6.6599(4) Å	$\beta = 104.014(3)^{\circ}$
c = 23.5921(16) Å	
$V = 3385.8(4) \text{ Å}^3$	
Absorption coefficient = $0.47 \text{ mm}^{-1}$	
F(000) = 1672	
Crystal dimensions (mm) =	= 0.38×0.32×0.29
$R[F^2 > 2\sigma(F^2)] = 0.04$	
$R_{\rm int} = 0.03$	
$R_w(F^2) = 0.127$	$-31 \le h \le 31$
$\theta$ range for data collection = 1.1 to 30.6 °	$-9 \le k \le 9$
Goodness to fit = $1.043$	$-31 \le l \le 33$
$\Delta \rho_{\rm max} = 1.48 \text{ e} \text{ Å}^{-3}$	
$\Delta \rho_{\rm min} = -1.04 \ e \ {\rm \AA}^{-3}$	
Data collection = Bruke	r SMART 1000
Absorption correction: SADABS	
No. of independent reflections = $10365$	
No. of reflections collected w	with $I > 2\sigma(I) = 8960$
No. of parameters = $496$	
Program used to solve the structural =	SHELXS-97
Program used to refine the stru-	uctural = SHELXL-97
CCDC Number: Crystallographic data for this struct Crystallographic Data Centre, CCDC number 778278. C	ture has been deposited with the Cambridge Copies of the data can be obtained free of charge

**Table 1.** Crystal and experimental data of the title compound

Crystallographic Data Centre, CCDC number 778278. Copies of the data can be obtained free of charge on application to the Director, CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (Fax: int. code (1223)336-033; e-mail for deposition: deposit@ccdc.cam.ac.uk).

Table 2. Atom	ic coordinate (×1	$(0^4)$ and eq	uivalent isotroj	pic displacement	t parameters (Å <sup>2</sup> ×	$10^3$ ) for non	hydrogen atoms

Atom	x	У	Ζ	$U_{(eq)}$
Mn1	0.245039 (9)	0.00309 (3)	-0.002465 (8)	0.01165 (7)
01	0.29406 (5)	-0.20288 (15)	-0.04521 (4)	0.01320 (18)
O1S	0.81976 (5)	0.22059 (16)	0.77135 (4)	0.0164 (2)
O2S	0.14066 (5)	0.47516 (16)	0.14632 (4)	0.0160 (2)
O2	0.36007 (5)	-0.46262 (17)	-0.03112 (4)	0.0160 (2)
O3S	0.17925 (5)	0.74071 (16)	-0.27215 (5)	0.0175 (2)
O4S	0.35518 (5)	-0.44881 (17)	-0.14720 (4)	0.0172 (2)
O5S	0.78473 (5)	-0.14113 (17)	0.80663 (5)	0.0217 (2)
05	0.19849 (5)	0.21276 (15)	0.04159 (4)	0.01255 (18)
O6S	0.21849 (6)	0.38719 (17)	-0.30782 (5)	0.0223 (2)
O6	0.13334 (5)	0.47412 (17)	0.02897 (5)	0.0174 (2)
N1	0.21675 (5)	0.25400 (17)	-0.06555 (5)	0.0112 (2)
N2	0.27593 (5)	-0.24192 (17)	0.06240 (5)	0.0111 (2)
N3	0.55638 (6)	0.08615 (18)	0.71707 (5)	0.0149 (2)
N4	0.45181 (6)	0.1743 (2)	0.97378 (6)	0.0199 (3)
N5	0.05762 (6)	0.10265 (19)	0.71429 (5)	0.0157 (2)
N6	-0.06763 (10)	0.1202 (3)	0.96002 (8)	0.0454 (6)
C1	0.32200 (6)	-0.3485 (2)	-0.01614 (5)	0.0107 (2)
C2	0.30915 (6)	-0.38610 (19)	0.04372 (5)	0.0100 (2)
C3	0.33099 (6)	-0.5549 (2)	0.07772 (6)	0.0108 (2)
C4	0.31877 (6)	-0.5698(2)	0.13287 (6)	0.0129 (2)
C5	0.28565 (6)	-0.4189 (2)	0.15217 (6)	0.0135 (2)
C6	0.26466 (6)	-0.2577 (2)	0.11551 (6)	0.0132 (2)
C8	0.17040 (6)	0.3591 (2)	0.01293 (5)	0.0108 (2)
C9	0.18278 (6)	0.39688 (19)	-0.04710 (5)	0.0102 (2)
C10	0.16108 (6)	0.5657 (2)	-0.08088(5)	0.0106 (2)
C11	0.17535 (6)	0.5847 (2)	-0.13509 (6)	0.0127 (2)
C12	0.21002 (6)	0.4372 (2)	-0.15381 (6)	0.0135 (2)
C13	0.22991 (6)	0.2737 (2)	-0.11770 (6)	0.0130 (2)
C15	0.59545 (7)	0.0820 (2)	0.77009 (6)	0.0154 (3)
C16	0.57629 (6)	0.0919 (2)	0.82189 (6)	0.0138 (2)
C17	0.51282 (6)	0.10539 (19)	0.81884 (6)	0.0114 (2)
C18	0.49061 (6)	0.1250 (2)	0.87291 (6)	0.0128 (2)
C19	0.52525 (7)	0.2389 (2)	0.91939 (6)	0.0168 (3)
C20	0.50484 (7)	0.2601 (2)	0.96950(7)	0.0192 (3)
C21	0.41674 (7)	0.0649 (2)	0.93055 (7)	0.0198 (3)
C22	0.43553 (7)	0.0367 (2)	0.87887 (7)	0.0167 (3)
C23	0.47159 (6)	0.1069 (2)	0.76389 (6)	0.0131 (2)
C24	0.49550(7)	0.0985 (2)	0.71458 (6)	0.0142 (2)
C25	0.09521 (7)	0.1030 (2)	0.76804 (6)	0.0160 (3)
C26	0.07416 (6)	0.1067 (2)	0.81890 (6)	0.0138 (2)
C27	0.01034 (6)	0.10728 (19)	0.81410 (6)	0.0109 (2)
C28	-0.01524 (6)	0.1138 (2)	0.86640 (6)	0.0132 (2)
C29	0.01074 (9)	0.2346 (2)	0.91424 (7)	0.0239 (3)
C30	-0.01746 (13)	0.2342 (3)	0.96132 (8)	0.0425 (6)
C31	-0.09281 (10)	0.0047 (3)	0.91502 (10)	0.0372 (5)
C32	-0.06763 (7)	-0.0007(2)	0.86758 (8)	0.0219 (3)
C33	-0.02932 (6)	0.1043 (2)	0.75832 (6)	0.0129 (2)
C34	-0.00373 (7)	0.1039 (2)	0.71009 (6)	0.0148 (2)
O10A	0.33815 (5)	-0.86557 (16)	0.03024 (5)	0.0161 (2)
C10A	0.36724 (6)	-0.7201 (2)	0.05748 (6)	0.0117 (2)
O12A	0.42553 (5)	-0.70586 (16)	0.07144 (4)	0.01464 (19)
09A	0.15081 (5)	0.87381 (16)	-0.03288 (5)	0.0165 (2)
C30A	0.12249 (6)	0.7260 (2)	-0.06154 (6)	0.0129 (2)
011A	0.06503 (5)	0.70878 (16)	-0.07629 (4)	0.0162 (2)

 Table 3. Selected bond distances (Å), and bond angels (°)

1401	e er bereeten bonna u	istunees (11), and oona ange	215 ( )
Mn1–O1	2.1485 (10)	Mn1–N1	2.2247 (12)
Mn1-O5	2.1493 (10)	Mn1-N2	2.2285 (11)
Mn1–O10A <sup>i</sup>	2.2056 (10)	Mn1–O9A <sup>ii</sup>	2.2156 (11)
O1-Mn1-O5	178.36 (4)	O10A <sup>i</sup> -Mn1-N1	91.44 (4)
O1-Mn1-O10A <sup>i</sup>	83.03 (4)	O9A <sup>ii</sup> -Mn1-N1	89.06 (4)
O5-Mn1-O10A <sup>i</sup>	95.35 (4)	O1-Mn1-N2	75.41 (4)
O1-Mn1-O9A <sup>ii</sup>	98.41 (4)	O5-Mn1-N2	104.35 (4)
O5-Mn1-O9A <sup>ii</sup>	83.21 (4)	O10A <sup>i</sup> -Mn1-N2	86.76 (4)
O10A <sup>i</sup> -Mn1-O9A <sup>ii</sup>	178.30 (4)	O9A <sup>ii</sup> -Mn1-N2	92.72 (4)
O1-Mn1-N1	104.99 (4)	N1-Mn1-N2	178.10 (4)
O5-Mn1-N1	75.20 (4)		

Symmetry codes: (i) x, y+1, z; (ii) x, y-1, z.

 Table 4. Hydrogen bond geometry (Å ,°)

D–H…A	d(D-H)	d(H···A)	d(D····A)	<(D–H…A)
O1S-H1B···O2S <sup>iii</sup>	0.85	1.96	2.7972 (15)	170
O1S-H1A···O5S	0.85	1.88	2.7231 (15)	173
O2S-H2B···N3 <sup>iv</sup>	0.85	2.06	2.8229 (15)	150
O2S-H2A···O6	0.85	1.89	2.7338 (14)	174
O3S-H3B···O4S <sup>v</sup>	0.85	1.95	2.7886 (15)	168
O3S-H3A···O6S	0.85	1.87	2.7141 (16)	171
O4S-H4B…N5 <sup>vi</sup>	0.85	2.06	2.8032 (16)	146
O4S-H4A···O2	0.85	1.87	2.7159 (14)	175
O5S-H5B…O1S <sup>vii</sup>	0.85	1.90	2.7479 (15)	175
O5S-H5A···O2S <sup>viii</sup>	0.85	1.99	2.8348 (16)	171
O6S-H6B···O3S <sup>ix</sup>	0.85	1.91	2.7558 (16)	177
O6S-H6A···O4S <sup>x</sup>	0.85	1.98	2.8230 (16)	170
N4-H4C…O12A <sup>xi</sup>	0.85	1.80	2.6339 (16)	166
N4-H4C…O10A <sup>xi</sup>	0.85	2.55	3.1378 (17)	127
O1S-H1B···O2S <sup>iii</sup>	0.85	1.96	2.7972 (15)	170
O1S-H1A···O5S	0.85	1.88	2.7231 (15)	173
O2S-H2B···N3 <sup>iv</sup>	0.85	2.06	2.8229 (15)	150
O2S-H2A…O6	0.85	1.89	2.7338 (14)	174
O3S-H3B···O4S <sup>v</sup>	0.85	1.95	2.7886 (15)	168
O3S-H3A···O6S	0.85	1.87	2.7141 (16)	171
O4S-H4B…N5 <sup>vi</sup>	0.85	2.06	2.8032 (16)	146
O4S-H4A···O2	0.85	1.87	2.7159 (14)	175
O5S-H5B···O1S <sup>vii</sup>	0.85	1.90	2.7479 (15)	175
O6S-H6B···O3S <sup>ix</sup>	0.85	1.91	2.7558 (16)	177
O6S—H6A···O4S <sup>x</sup>	0.85	1.98	2.8230 (16)	170
C13—H13…O6S <sup>ix</sup>	0.93	2.58	3.4644 (18)	160
C20—H20…O2 <sup>viii</sup>	0.93	2.56	3.2897 (19)	135
C21—H21…O1 <sup>xii</sup>	0.93	2.50	3.4185 (19)	169
C24—H24…O11A <sup>vi</sup>	0.93	2.44	3.2965 (17)	153
C31—H31…O5 <sup>xiii</sup>	0.93	2.27	3.133 (2)	154
C34—H34···O12A <sup>xiv</sup>	0.93	2.45	3.3377 (17)	159

Symmetry codes: (iii) -x+1, -y+1, -z+1; (iv) x-1/2, -y+1/2, z-1/2; (v) -x+1/2, y+3/2, -z-1/2; (vi) -x+1/2, y-1/2, -z+1/2; (vii) -x+3/2, y-1/2, -z+3/2; (viii) -x+1, -y, -z+1; (ix) -x+1/2, y-1/2, -z-1/2; (x) -x+1/2, y+1/2, -z-1/2; (xi) x, y+1, z+1; (xii) x, y, z+1; (xiii) -x, -y, -z+1; (xiv) x-3/2, -y-3/2, z-1/2.

According to the crystal structure of the compound, the central atom is attached to two (py- $(2,3-dc)^{2-}$  ligands through their N and O atoms in a planar square arrangement. Also, pyridine-2,3dicarboxylate ligand,  $(py-2,3-dc)^{2-}$ , plays a bridging role. Its two carboxylic groups adopt monodentate coordination mode, one together with pyridine N atom coordinated to one Mn(III) ion, the other coordinates to second equivalent Mn(III) atom, forming polymeric chains.

In the compound, O9A and O10A atoms from bridging (py-2,3-dc)<sup>2-</sup> ligands [O10A<sup>i</sup>-Mn1-O9A<sup>ii</sup>  $= 178.30 (4)^{\circ}$ ; i: x, y + 1, z, ii: x, y-1, z], and O1, O5, N1 and N2 atoms [N1-Mn1-N2 = 178.10(4)° and O1-Mn1-O5 =  $178.36(4)^{\circ}$ ] are attached to the central atom. Therefore, the geometry of the resulting MnN<sub>2</sub>O<sub>4</sub> coordination fragment can be described as slightly distorted octahedral.

 $Mn1-O9^{ii}$ and  $Mn1-O10^{i}$ The bonds [2.2156(11) and 2.2056(10) Å] are significantly longer than other Mn-O bonds [2.1485(10) and 2.1493(10) Å]. Also, the Mn–O bonds are almost perpendicular to the square plane and by connecting Mn (III) ions together result in the formation of polymeric chains along b direction. The metal-ligand bond distances are in agreement with published Mn (III) complexes [17,18]. There are also six uncoordinated water molecules, one neutral 4,4'-bipy molecule and one

monoprotonated  $(4,4'-bipyH)^+$  cation as counterion.

The two pyridine rings in the neutral 4,4'-bipy molecule and  $(4,4'-bipyH)^+$  cation are twisted with respect to each other, making dihedral angles of 40.69(8) and 37.01(6)°, respectively.

A noticeable feature of the title compound is the presence of C=O $\cdot\cdot\cdot\pi$  stacking interactions between C=O groups of carboxylate groups with aromatic rings of (py-2,3-dc) - and (4,4'-bipyH) + units. The C=O··· $\pi$  distances (measured to the center of phenyl ring) are 3.6986(12) Å for C1=O2…Cg1 (x, -1+y, z), 3.7017(13) Å for C8=O6…Cg2 (x, 1+y, z) and 3.7343(13) Å for C10A=O12A...Cg3 (1-x, -1y, 1-z), [Cg1, Cg2 and Cg3 are the centroids for N1/C9-C13, N2/C2-C6 and N4/C18-C22 rings, respectively]. There is also an interesting C-H $\cdots\pi$ interaction between C29-H29-Cg3 and phenyl ring of  $(py-2,3-dc)^{2-}$  unit with H··· $\pi$  distance of 2.94 Å (x, y, 1+z) (Fig. 2). Also a considerable  $\pi - \pi$ stacking interactions between aromatic rings of 4,4'-bipy and  $(4,4'-bipyH)^+$ fragments with distances of 3.6065(8) and 3.7216(8) Å for Cg4…Cg5 (1/2-x, -1/2+y, 3/2-z) and (1/2-x, 1/2+y, 3/2-z) are observed in the prepared compound [Cg4 and Cg5 are centroids for N3/C15-C17,C23,C24 and N5/C25-C27,C33,C34 rings, respectively] (Fig.3) [19-23].



3 702

64



**Fig. 3**  $\pi$ - $\pi$  stacking interactions between aromatic rings of 4,4'-bipy and (4,4'-bipyH)<sup>+</sup> fragments with distances 3.6065(8) and 3.7216(8) Å for Cg4…Cg5 (1/2-x, -1/2 + y, 3/2-z) and (1/2-x, 1/2+y, 3/2-z) [Cg4 and Cg5 are the centroids for N3/C15-C17,C23,C24 and N5/C25-C27,C33,C34 rings, respectively].

As shown in Figure 4, in the crystal structure, a wide range of hydrogen bonding interactions consisting of O–H···O, O–H···N, N–H···O and C–H···O types with D···A ranging from 2.6339(16) to 3.4644(18) Å connect the different parts together to form layers in which the space between anionic chains is filled with cations and uncoordinated water molecules.

In addition, hydrogen bonds between water molecules create  $(H_2O)_n$  clusters and these units connect the cations and anions together throughout the structure. These cyclic  $(H_2O)_n$  clusters are shown at Figure 5 and are located in the space between layers of cations and anions. The crystal packing diagram of the polymeric compound is illustrated in Figure 6.



Fig. 4 Cyclic pentamer water clusters formed by distinct intermolecular O-H…O and hydrogen bonds.



**Fig. 5** Hydrogen bonding between different species, the cyclic pentamer clusters brings together the cations and anions. Hydrogen bonds are shown as dashed lines.



Fig. 6 Crystal packing diagram of the title compound. Hydrogen bonds are shown by dashed lines.

## References

[1] Yi L., Din B., Zhao B., Cheng P., Liao D. -Z., Yan S. -P., Jiang Z. -H., "Novel triazole-bridged cadmium coordination polymers varying from zero- to three-dimensionality", Inorganic Chemistry, 43 (2004) 33-43.

[2] Withersby M. A., Blake A. J., Champness N. R., Cooke P. A., Hubberstey P., Li W. S., Schroder

M., "Solvent control in the synthesis of 3,6-bis (pyridin-3-yl) - 1,2,4,5 – tetrazine - bridged cadmium (II) and zinc(II) coordination polymers", Inorganic Chemistry, 38 (1999) 2259-2266.

[3] Shin D. M., Lee I. S., Chung Y. K., Lah M. S., "Synthesis and characterization of novel grid coordination polymer networks generated from *unsymmetrically bridging ligands",* Inorganic Chemistry, 42 (2003) 5459-5461.

[4] Hagrman D., Haushalter R. P., Zubieta J., "Organic/Inorganic composite materials: Hydrothermal syntheses and structures of the one-, and three-dimensional copper(II) two-. sulfate-organodiamine phases  $[Cu(H_2O)_3(4,4)]$ *bipyridine*)( $SO_4$ )]·2 $H_2O_5$ ,  $[Cu(bpe)_2]$  $[Cu(bpe)(H_2O)_2(SO_4)_2] \cdot 2H_2O,$ and  $[Cu(bpe)(H_2O)(SO_4)]$  (bpe = trans-1,2-Bis(4pyridyl)ethylene)", Chemistry of Materials, 10 (1998) 2091-20100.

[5] Reineke T. M., Eddaoudi M., Molar D., Keeffe M. O, Yaghi O. M., "Large free volume in interpenetrating networks: The role of secondary building units exemplified by  $Tb_2(ADB)_3[(CH_3)_2SO]_4 \cdot 16[(CH_3)_2SO]"$ , Journal of American Chemical Society, 122 (2000) 4843-4844.

[6] Sharma C. V. K., Rogers R. D., "Perspectives of crystal engineering", Materials Today, 1(3) (1998) 27-30.

[7] Seo J. S., Whang D., Lee H., Jun S. I., Oh J., Jeon Y. J., Kim K., "A homochiral metal-organic porous material for enantioselective separation and catalysis", Nature, 404 (2000) 982-986.

[8] H. B. Xu, X. F. Li, D. Yan, "Crystal structure and photo- and electroluminescent properties of a sandglass terbium cluster", Inorganic Chemistry Communications, 11 (2008) 1187-1189.

[9] Deng H., Li Y. H., Qiu Y. C., Liu Z. H., Zeller M., "Construction of isostructural Ln-Ag (Ln = Eu; Tb; Dy) nano-channel heterometallic coordination frameworks based on pyrazine-2-carboxylate and oxalate ligands", Inorganic Chemistry Communications, 11 (2008) 1151-1154.

[10] Ma L. F., Wang L. Y., Wang J. G., Wang Y. F., Feng X., "Syntheses, structure and properties of *Zn(II) and Co(II) complexes of N-benzesulfonyl-1-glutamic acid ligand*", Inorganica Chimica Acta, 359 (2006) 2241-2245.

[11] Rueff J. -M., Masciocchi N., Rabu P., Sironi A., Skoulios A., "Structure and magnetism of a polycrystalline transition-metal soap:  $Co^{II}[OOC(CH_2)_{10}COO](H_2O)_2$ ", European Journal of Inorganic Chemistry, 2001 (2001), 2843-2848. [12] Blake A. J., Champness N. R., Crew M., Hanton L. R., Parsons S., Schroder M., "A new Cu<sup>l</sup>(SCN) structural motif: Synthesis of an uncharged three-dimentional coordination network", Journal of the Chemical Society, Dalton Transactions, (1998) 1533-1534.

[13] Maji T. K., Sain S., Mostafa G., Lu T. H., Ribas J., Monfort M., Chaudhuri N. R., "Magneto-structural correlations in 2D and 3D extended structures of manganese(II)-malonate systems", Inorganic Chemistry, 42 (2003) 709-716.
[14] Rather B., Zaworotko M., "A 3D metalorganic network, [Cu<sub>2</sub>(glutarate)<sub>2</sub>(4,4'bipyridine)], that exhibits single-crystal to singlecrystal dehydration and rehydration", Chemical

[15] Liu T. -F., Sun H. -L., Gao S., Zhang S. -W., Lau T. -C., "Ferromagnetic ordering and metamagnetism in malonate bridged 3D diamondlike and honey-like networks:  $[Cu(mal)(MDF)]_n$ and  $\{[Cu(mal)(0.5pyz)].nH_2O\}$  (mal=malonate dianion, MDF=N,N-dimethylformamide, pyz=pyrazin ", Inorganic Chemistry, 42 (2003) 4792-4794.

Communication, 7 (2003) 830-831.

[16] Sheldrick G. M., "A short history of SHELX", Acta Crystallographica Section A, 64 (2008) 112-122.

[17] Tynan, E., Jensen, P., Kruger, P. E., Lees, A. C., "Solvent templated synthesis of metalorganic frameworks: structural characterization and properties of the 3D network isomers  $\{[Mn(dcbp)]\cdot/_2DMF\}_n$  and  $\{[Mn(dcbp)]\cdot2H_2O\}_n$ ", Chemical Communications, (2004) 776-777.

[18] Marioni P.-A., Marty W., Stoeckli-Evans H., Whitaker C., "Coordination polymers of *Mn(III) with the ligand pyrazine-2, 3, 5, 6-tetracarboxylic acid* ", Inorganica chimica Acta, 219 (1994) 161-168.

[19] Soleimannejad J., Attar Gharamaleki J., Aghabozorg H., Mohammadzadeh Azar Golenji Y., "4,4'-Bipyridinium bis(μ-4-oxo-1,4dihydropyridine-2,6-dicarboxylato) bis [aquahydroxidoantimonate(III)] dihydrate ", Acta Crystallographica Section E, 66 (2010) m467m468.

[20] Soleimannejad J., Hooshmand H., Ghadermazi M., Attar Gharamaleki J., "The monoclinic polymorph of catenapoly[[diaquamanganese(II)]-µ-oxalato-

 $\kappa^4 O^1, O^2: O^{1'}, O^{2'} J''$ , Acta Crystallographica Section E, 63 (2010) m2389-m2390.

[21] Soleimannejad J., Aghabozorg H., Morsali A., Hemmati F., Manteghi F., "4,4'-*Bipyridinium bis(2-carboxypyridine-3carboxylate)* ", Acta Crystallographica Section E, 65(1) (2009) 0153-0153. [22] Soleimannejad J., Aghabozorg H., Hooshmand S., Adams H., "4,4'-Bipyridinediium triaquabis(pyridine-2,6-dicarboxylato) strontium (II) trihydrate", Acta Crystallographica Section E, 63(12) (2007) m3089-m3090.

[23] Soleimannejad J., Nasibipoor M., Hemmati F., "*Crystal structure of a Cr(III) complex containing nitrilotriacetic acid and 4,4'-bipyridine*", Iranian Journal of Crystallography and Mineralogy, 20(1) (2012) 1-8.